
WAINUA is a prescription medication for the treatment of polyneuropathy caused by hereditary transthyretin-mediated amyloidosis (hATTR-PN) in adults.





Individual results may vary.

*As measured by mNIS+7 score

IMPORTANT SAFETY INFORMATION

WAINUA may cause side effects, including:

Low vitamin A level

See pages 2-3 for additional information.

Please see additional Important Safety Information on pages 2 and 3 and full <u>Prescribing Information</u>, including Patient Information, for WAINUA.



IMPORTANT SAFETY INFORMATION

WAINUA may cause side effects, including:

• Low vitamin A level, which is a serious, but common side effect of treatment with WAINUA. Your doctor should tell you to take vitamin A supplements. Do not take more than the amount of vitamin A your doctor has recommended. Call your doctor if you develop eye problems such as difficulty seeing at night or in low-lit areas (night blindness), or dry eyes. If you develop eye problems, your doctor should refer you to an eye doctor

The most common side effects include decreased vitamin A and vomiting.

Before you take WAINUA, tell your healthcare provider:

• if you are pregnant or breastfeeding, or plan to become pregnant or breastfeed. It is not known if WAINUA can harm your baby. Changes in vitamin A levels and vitamin A supplementation related to use of WAINUA may harm your unborn baby. It is not known if WAINUA passes into breast milk


 about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements.
 Especially tell your healthcare provider if you take vitamin A or beta-carotene supplements

You may <u>report side effects related to AstraZeneca products</u>. ☑

No additional laboratory monitoring is needed when you take WAINUA.

Please see full <u>Prescribing</u>
<u>Information</u>, including <u>Patient</u>
Information, for WAINUA.





IMPORTANT SAFETY INFORMATION (continued)

WAINUA may cause side effects, including:

• Low vitamin A level, which is a serious, but common side effect of treatment with WAINUA. Your doctor should tell you to take vitamin A supplements. Do not take more than the amount of vitamin A your doctor has recommended. Call your doctor if you develop eye problems such as difficulty seeing at night or in low-lit areas (night blindness), or dry eyes. If you develop eye problems, your doctor should refer you to an eye doctor





IMPORTANT SAFETY INFORMATION (continued)

The most common side effects include decreased vitamin A and vomiting.



In adults with hATTR-PN, this is a chance to

Stop hATTR neuropathy progression.*

Start improving your quality of life.*




WAINUA can prevent your neuropathy from getting worse **so you may feel stronger.**



WAINUA can improve quality of life, allowing you to do more of the things you want to do.

Individual results may vary.



*As measured by mNIS+7 score and Norfolk QoL-DN score
'Improvement was defined as the difference between the change from baseline in the WAINUA group and a placebo group from a similar study.

The mNIS+7 scale assessed nerve function by measuring things such as muscle strength, sensation, and reflexes.

The Norfolk QoL-DN questionnaire assessed patients' perceptions of their neuropathy symptoms and its impact on daily activities such as dressing, bathing, or walking.

IMPORTANT SAFETY INFORMATION (continued)

WAINUA may cause side effects, including:

• Low vitamin A level, which is a serious, but common side effect of treatment with WAINUA. Your doctor should tell you to take vitamin A supplements. Do not take more than the amount of vitamin A your doctor has recommended. Call your doctor if you develop eye problems such as difficulty seeing at night or in low-lit areas (night blindness), or dry eyes. If you develop eye problems, your doctor should refer you to an eye doctor

The most common side effects include decreased vitamin A and vomiting.



WAINUA puts control of administration in your hands.



Take WAINUA at home or on the go

WAINUA is the only autoinjector for hATTR-PN. Make sure you store it properly (see below).



It's taken once monthly

You can take WAINUA by yourself or someone can help you.



The actual injection takes just 10 seconds

Be sure to read the complete Instructions for Use.



Store at room temperature*

The WAINUA autoinjector arrives in a cooler; store it in its carton in a refrigerator. If needed, WAINUA can be kept at room temperature in the original carton for up to 6 weeks.

*If needed, WAINUA can be kept at room temperature up to 86 °F (30 °C). Do not let WAINUA reach temperatures above 86 °F (30 °C) or expose to heat. Throw away WAINUA if kept at room temperature longer than 6 weeks. 'Store in its original carton between 36 °F to 46 °F (2 °C to 8 °C). Do not freeze.

IMPORTANT SAFETY INFORMATION (continued)

WAINUA may cause side effects, including:

 Low vitamin A level, which is a serious, but common side effect of treatment with WAINUA. Your doctor should tell you to take vitamin A supplements. Do not take more than the amount of vitamin A your doctor has recommended. Call your doctor if you develop eye problems such as difficulty seeing at night or in low-lit areas (night blindness), or dry eyes. If you develop eye problems, your doctor should refer you to an eye doctor

Please see additional Important Safety Information on pages 2 and 3.



Before taking WAINUA, keep in mind:

- WAINUA suppresses the production of the protein that transports vitamin A and, therefore, vitamin A supplementation is recommended. See Important Safety Information below
- Your healthcare provider should demonstrate how to use the autoiniector



Scan the QR code to watch a step-by-step instructional video

or visit WAINUA.com/HowToVideo.



WAINUA Way is committed to supporting you

We know you could handle this on your own. But with the WAINUA WAY support program, you don't have to.



Savings and affordability options

AstraZeneca Access 360[™] can answer questions about insurance, out-of-pocket costs, and assistance programs.



A dedicated point of contact

Your Patient Education Manager will share resources on how to take WAINUA, give you important updates, and more.



Educational resources

You'll get information about WAINUA and resources to help get you started with treatment.

To Learn More or to Enroll in WAINUA Way



Scan this QR code, visit WAINUAway.com, or call 1-844-2-WAINUA (844-292-4682).






Tell your story. Inspire people like you.

The AZ US Patient Voices program is an opportunity to share your unique experiences with WAINUA. Email us at MyStory@AstraZeneca.com to learn more.









Ask your healthcare provide if WAINUA may be the right treatment for you.

Scan this QR Code or **visit WAINUA.com**/Results.

Please see Important Safety Information on pages 2 and 3.



WAINUA is a trademark of the AstraZeneca group of companies. ©2024 AstraZeneca. All rights reserved. US-92244 Last Updated 9/24

