

This tool is to help you keep track of any signs or symptoms you are experiencing that may be related to polyneuropathy caused by hereditary ATTR amyloidosis (hATTR-PN).

Symptoms of hATTR-PN are not the same for everyone and may seem unrelated. Use this tracker to write down when you experience symptoms, how severe they are, and how they impact your daily life. Sharing this with your doctor can help them better understand your health.

Review the Important Safety Information for WAINUA on the following page and discuss any side effects that bother you or do not go away with your healthcare team.

This symptom tracker is not a substitute for reporting symptoms to your healthcare team. Report any symptoms that concern you to your healthcare team right away.

1	2	3	4	5
START HERE Fill out in 5 easy steps	Select your hATTR-PN symptoms	Indicate the severity on a scale from 1 to 5, with 5 being severe	Since your last visit with your doctor, indicate if your symptoms have: I - Improved S - Stayed the same W - Worsened	Write down the dates you've experienced these symptoms
Indicate the impact on daily life: A) Difficulty with daily activities, including work B) Reduced social interaction C) Reduced physical activity (eg, walking, lifting) D) Reduced emotional well-being				
☒ Example symptom	1 2 ☒ 4 5	I ☒ W	6/3-7/3	A ☒ C D

hATTR-PN SYMPTOMS IN THE HANDS, ARMS, LEGS, AND FEET

☐ Tingling or pain in your hands or feet	1 2 3 4 5	I S W		A B C D
☐ Aching pain in both hands and/or forearms	1 2 3 4 5	I S W		A B C D
☐ Reduced grip	1 2 3 4 5	I S W		A B C D
☐ Loss of sensitivity to touch or temperature	1 2 3 4 5	I S W		A B C D
☐ Numbness or weakness in the legs and feet	1 2 3 4 5	I S W		A B C D
☐ Difficulty walking or moving around	1 2 3 4 5	I S W		A B C D

hATTR-PN SYMPTOMS RELATED TO THE HEART, BLOOD VESSELS, AND CIRCULATION

☐ Abnormal heartbeat	1 2 3 4 5	I S W		A B C D
☐ Shortness of breath	1 2 3 4 5	I S W		A B C D
☐ Fatigue	1 2 3 4 5	I S W		A B C D
☐ Swelling in the feet or ankles	1 2 3 4 5	I S W		A B C D
☐ Dizziness upon standing	1 2 3 4 5	I S W		A B C D
☐ Lightheadedness	1 2 3 4 5	I S W		A B C D

Please see Approved Use and Important Safety Information on the next page.

1	2	3	4	5
START HERE Fill out in 5 easy steps	Select your hATTR-PN symptoms	Indicate the severity on a scale from 1 to 5, with 5 being severe	Since your last visit with your doctor, indicate if your symptoms have: I - Improved S - Stayed the same W - Worsened	Write down the dates you've experienced these symptoms
				Indicate the impact on daily life : A) Difficulty with daily activities, including work B) Reduced social interaction C) Reduced physical activity (eg, walking, lifting) D) Reduced emotional well-being

☒ Example symptom	1 2 ☒ 4 5	I ☒ W	6/3-7/3	A ☒ C D
---	---	---	---------	---

hATTR-PN SYMPTOMS RELATED TO DIGESTION, NUTRITION, AND WEIGHT

☐ Diarrhea	1 2 3 4 5	I S W		A B C D
☐ Constipation	1 2 3 4 5	I S W		A B C D
☐ Loss of bowel or bladder control	1 2 3 4 5	I S W		A B C D
☐ Reduced appetite	1 2 3 4 5	I S W		A B C D
☐ Unintended weight loss	1 2 3 4 5	I S W		A B C D

OTHER hATTR-PN SYMPTOMS

☐ Blurry vision	1 2 3 4 5	I S W		A B C D
☐ Eye floaters	1 2 3 4 5	I S W		A B C D
☐ Sexual dysfunction	1 2 3 4 5	I S W		A B C D
☐ Anxiety and/or depression	1 2 3 4 5	I S W		A B C D

Approved Uses

WAINUA injection, for subcutaneous use, 45 mg is a prescription medicine used to treat adults with polyneuropathy of hereditary transthyretin-mediated amyloidosis (hATTR-PN).

Important Safety Information

WAINUA may cause side effects, including:

- **Low vitamin A level**, which is a serious, but common side effect of treatment with WAINUA. Your doctor should tell you to take vitamin A supplements. **Do not** take more than the amount of vitamin A your doctor has recommended. Call your doctor if you develop eye problems such as difficulty seeing at night or in low-lit areas (night blindness), or dry eyes. If you develop eye problems, your doctor should refer you to an eye doctor

The most common side effects include decreased vitamin A and vomiting.

Before you take WAINUA, tell your healthcare provider:

- if you are pregnant or breastfeeding, or plan to become pregnant or breastfeed. It is not known if WAINUA can harm your baby. Changes in vitamin A levels and vitamin A supplementation related to use of WAINUA may harm your unborn baby. It is not known if WAINUA passes into breast milk
- about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take vitamin A or beta-carotene supplements

Please read full [Prescribing Information](#), including [Patient Information](#), for WAINUA.

You may [report side effects related to AstraZeneca products](#) .